

LETTERS

cases can be expected as pressure on marine fish stocks and consumption of uncooked fish increase. As a protective measure against this emerging foodborne zoonotic threat, the public should be made aware of risks associated with consumption of fresh, raw marine fish.

Acknowledgments

We thank Amanda Rogers from SA Pathology for her contribution.

References

1. Kuchta R, Serrano-Martínez ME, Scholz T. Pacific broad tapeworm *Adenocephalus pacificus* as a causative agent of globally reemerging diphyllbothriosis. *Emerg Infect Dis*. 2015;21:1697–703. <http://dx.doi.org/10.3201/eid2110.150516>
2. Hernández-Orts JS, Scholz T, Brabec J, Kuzmina T, Kuchta R. High morphological plasticity and global geographical distribution of the Pacific broad tapeworm *Adenocephalus pacificus* (syn. *Diphyllbothrium pacificum*): molecular and morphological survey. *Acta Trop*. 2015;149:168–78. <http://dx.doi.org/10.1016/j.actatropica.2015.05.017>
3. Johnston TH. Entozoa from the Australian hair seal. *Proc Linn Soc N S W*. 1937. [cited 2016 Jan 20]. <http://www.biodiversitylibrary.org/page/34903721#page/59/mode/1up>
4. Wicht B, Yanagida T, Scholz T, Ito A, Jiménez JA, Brabec J. Multiplex PCR for differential identification of broad tapeworms (Cestoda: *Diphyllbothrium*) infecting humans. *J Clin Microbiol*. 2010;48:3111–6. <http://dx.doi.org/10.1128/JCM.00445-10>
5. Mariaux J. A molecular phylogeny of the Eucestoda. *J Parasitol*. 1998;84:114–24. <http://dx.doi.org/10.2307/3284540>
6. Luton K, Walker D, Blair D. Comparisons of ribosomal internal transcribed spacers from two congeneric species of flukes (Platyhelminthes: Trematoda: Digenea). *Mol Biochem Parasitol*. 1992;56:323–7. [http://dx.doi.org/10.1016/0166-6851\(92\)90181-I](http://dx.doi.org/10.1016/0166-6851(92)90181-I)
7. Pastor-Valle J, González LM, Martín-Clemente JP, Merino FJ, Gottstein B, Gárate T. Molecular diagnosis of diphyllbothriasis in Spain, most presumably acquired via imported fish, or sojourn abroad. *New Microbes New Infect*. 2014;2:1–6. <http://dx.doi.org/10.1002/2052-2975.28>
8. Kuchta R, Esteban JG, Brabec J, Scholz T. Misidentification of *Diphyllbothrium* species related to global fish trade, Europe. *Emerg Infect Dis*. 2014;20:1955–7. <http://dx.doi.org/10.3201/eid2011.140996>
9. Anantanawat S, Kiermeier A, McLeod C, Sumner J. South Australian Research and Development Institute. A semi-quantitative risk assessment of harmful parasites in Australian finfish. April 2012 [cited 2016 Jan 20]. <http://safefish.com.au/wp-content/uploads/2013/03/Semi-quantitative-risk-assessment-of-Harmful-Parasites.pdf>
10. New South Wales Food Authority. Seafood safety scheme—periodic review of the risk assessment. 2012. [cited 2016 Jan 20]. http://foodauthority.nsw.gov.au/_Documents/scienceandtechnical/seafood_safety_scheme.pdf

Address for correspondence: Gerhard F. Weldhagen, Microbiology and Infectious Diseases Laboratory, SA Pathology, PO Box 14, Rundle Mall, Adelaide, SA5000, Australia; email: gerhard.weldhagen@sa.gov.au

Influenza A(H7N7) Virus among Poultry Workers, Italy, 2013

Simona Puzelli,¹ Caterina Rizzo,¹ Concetta Fabiani, Marzia Facchini, Paolo Gaibani, Maria P. Landini, Carlo Gagliotti, Maria L. Moro, Roberto Rangoni, Luisa Loli Piccolomini, Alba C. Finarelli, Marco Tamba, Giovanni Rezza, Silvia Declich, Isabella Donatelli, Maria R. Castrucci

Author affiliations: Istituto Superiore Sanità, Rome, Italy (S. Puzelli, C. Rizzo, C. Fabiani, M. Facchini, G. Rezza, S. Declich, I. Donatelli, M.R. Castrucci); St. Orsola University Hospital, Bologna, Italy (P. Gaibani, M.P. Landini); Agenzia Sanitaria Regionale Emilia-Romagna Region, Bologna (C. Gagliotti, M.L. Moro, R. Rangoni, L.L. Piccolomini, A.C. Finarelli); Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna, Brescia, Italy (M. Tamba)

DOI: <http://dx.doi.org/10.3201/eid2208.160246>

To the Editor: In August 2013, an outbreak of infection with highly pathogenic influenza A(H7N7) virus occurred in Emilia-Romagna, Italy, and >1 million birds were culled (1). Prevention measures were immediately applied, and all workers involved in culling activities wore personal protective equipment (PPE), including face masks with eye protection. These workers were monitored for clinical symptoms, and 3 workers with laboratory-confirmed cases of conjunctivitis caused by infection with influenza A(H7N7) virus were reported during the 3-week outbreak (2). Workers did not receive chemoprophylaxis.

A serologic study was conducted in December 2013 to identify potential asymptomatic infections following exposure to influenza A(H7N7) virus. This study was approved by the ethics committee of the Istituto Superiore di Sanità (protocol no. PRE787/13CE13/401).

A total of 93 of 140 workers directly involved in culling activities, including the 3 confirmed case-patients with conjunctivitis, participated in the study. All participants completed a questionnaire that obtained information for demographics, poultry exposure, and use of PPE.

Paired acute-phase and convalescent-phase serum samples were available only for the 3 H7 subtype-positive persons with conjunctivitis. We tested these paired serum samples and single serum samples obtained from virus-exposed workers for antibodies against influenza A(H7N7) virus strain A/Italy/3/2013 (2) by using hemagglutination inhibition (HI) and microneutralization (MN) assays (3,4).

¹These authors contributed equally to this article.

Table. HI and MN antibody titers for influenza A(H7N7) virus and other H7 subtype viruses in serum samples of 5 men, Italy, 2013*

Person ID and phase type†	Age, y	Activity of person	Date of sample collection	Virus strain (subtype) and titer							
				A/It/3/2013 (H7N7)		A/Tk/It/3889/1999 (H7N1)		A/Tk/It/214845/2002 (H7N3)		A/Ck/It/2837-54/2007 (H7N3)	
				HI	MN	HI	MN	HI	MN	HI	MN
1											
Acute	51	PW, culling	Sep 6	10	<10	<10	<10	<10	<10	<10	<10
Convalescent			Dec 6	20	35	<10	<10	<10	<10	<10	<10
2											
Acute	46	Culling	Sep 6	10	<10	<10	<10	<10	<10	<10	<10
Convalescent			Dec 11	20	62	<10	<10	<10	<10	<10	<10
3											
Acute	49	Culling	Sep 7	<10	<10	<10	NT	<10	NT	<10	NT
Convalescent			Dec 23	10	87	<10	<10	<10	<10	<10	<10
FO10‡	34	Culling	Dec 23	20	72	<10	<10	<10	<10	<10	<10
RA32‡	55	PW, culling	Dec 11	20	33	<10	<10	<10	<10	<10	<10

*Bold indicates titers of seropositive persons (HI positive results confirmed 3 times by MN). Values for 1 of 3 MN assays that showed similar results are reported. Seropositive persons were selected from 93 persons who participated in the study among 140 persons involved in culling activities.

HI, hemagglutination inhibition; ID, identification; MN, microneutralization; NT, not tested; PW, poultry worker.

†Persons 1, 2, and 3 had laboratory-confirmed cases of conjunctivitis caused by infection with influenza A(H7N7) virus.

‡Asymptomatic person.

Other H7 subtype viruses previously circulating in Italy were included in the analysis to rule out potential cross-reactivity with influenza A(H7N7) virus (5). HI titers ≥ 10 and MN titers ≥ 20 were considered positive; only HI-positive serum samples confirmed 3 times by MN assay were considered positive results for influenza A(H7N7) virus.

We detected antibodies against influenza A(H7N7) virus in convalescent-phase serum samples from the 3 H7 subtype-positive patients and 2 asymptomatic persons but found no seropositivity against other H7 subtype viruses (Table). Because of lack of acute-phase serum samples, we could not assess whether seropositivity for the 2 asymptomatic persons, 1 (RA32) of whom worked with poultry before the outbreak, was caused by infection acquired during the outbreak. All workers were trained and most participants, including the 2 asymptomatic influenza A(H7N7) virus-seropositive persons, reported that PPE was commonly used during culling on infected premises. Nevertheless, it is likely that worker compliance with PPE was not always 100% during the 3-week outbreak because of poor knowledge and real perception of biologic risks among workers.

Future efforts should ensure timely collection of paired serum samples from all workers involved in avian influenza outbreaks, especially when infections occur in humans. Strict compliance with recommended preventive control measures and serologic surveillance programs are crucial to avoid and eventually assess risk for infections with avian influenza viruses in persons exposed to infected poultry.

This study was supported by the Italian Ministry of Health (grants CCM 2012 and RF-2010-2318269).

References

- Bonfanti L, Monne I, Tamba M, Santucci U, Massi P, Patreggiani T, et al. Highly pathogenic H7N7 avian influenza in Italy. *Vet Rec*. 2014;174:382. <http://dx.doi.org/10.1136/vr.102202>
- Puzelli S, Rossini G, Facchini M, Vaccari G, Di Trani L, Di Martino A, et al. Human infection with highly pathogenic A(H7N7) avian influenza virus, Italy, 2013. *Emerg Infect Dis*. 2014;20:1745–9. <http://dx.doi.org/10.3201/eid2010.140512>
- Centers for Disease Control and Prevention. Modified hemagglutination-inhibition (HI) assay using horse RBCs for serologic detection of antibodies to H7 subtype avian influenza virus in human sera. Version 1 [cited 2016 April 8]. https://consise.tghn.org/site_media/media/articles/160713_Modified_Hemagglutination_Inhibition_Assay_Using_Horse_RBCs.pdf
- World Health Organization. Serological detection of avian influenza A(H7N9) infections by microneutralization assay. Version 23, May 2013 [cited 2016 April 8]. http://www.who.int/influenza/gisrs_laboratory/cnic_serological_diagnosis_microneutralization_a_h7n9.pdf?ua=1
- Puzelli S, Di Trani L, Fabiani C, Campitelli L, De Marco MA, Capua I, et al. Serological analysis of serum samples from humans exposed to avian H7 influenza viruses in Italy between 1999 and 2003. *J Infect Dis*. 2005;192:1318–22. <http://dx.doi.org/10.1086/444390>

Address for correspondence: Simona Puzelli, National Influenza Centre, Department of Infectious, Parasitic and Immune-Mediated Diseases, Istituto Superiore di Sanità, Viale Regina Elena, 299-00161, Rome, Italy; email: simona.puzelli@iss.it

Increase in Eyeworm Infections in Eastern Europe

Vito Colella, Zvezdelina Kirkova, Éva Fok, Andrei D. Mihalca, Suzana Tasić-Otašević, Adnan Hodžić, Filipe Dantas-Torres, Domenico Otranto

Author affiliations: Università Degli Studi di Bari, Bari, Italy (V. Colella, F. Dantas-Torres, D. Otranto); Trakia University, Stara Zagora, Bulgaria (Z. Kirkova); Szent István University, Budapest, Hungary (É. Fok); University of Agricultural Sciences and